

The origin of the highest energy cosmic rays

Do all roads lead back to Virgo?

Eun-Joo Ahn¹, Gustavo Medina-Tanco², Peter L. Biermann³, and Todor Stanev⁴

¹*Department of Physics, Seoul National University, Seoul, Korea*

²*Institute of Astronomy and Geophysics, University of Sao Paulo, Sao Paulo, Brasil*

³*Max Planck Institut für Radioastronomie, Bonn, Germany*

⁴*Bartol Research Institute, University of Delaware, Newark, DE, USA*

Introducing a simple Galactic wind model patterned after the solar wind we show that back-tracing the orbits of the highest energy cosmic events suggests that they may all come from the Virgo cluster, and so probably from the active radio galaxy M87. This confirms a long standing prediction, as well as a theoretical model for the radio galaxy jet emission of M87. With this picture in hand, one clear expectation is that those powerful radio galaxies that have their relativistic jets stuck in the interstellar medium of the host galaxy, such as 3C147, will yield limits on the production of any new kind of particle, expected in some extensions of the standard model in particle physics.

PACS numbers: 98.70.Sa, 98.35.Gi, 78.35.Nq, 78.35.Eg

I. INTRODUCTION

The origin of the highest energy particles observed in the universe continues to present a major enigma to physics. These particles reach energies as high as $3 \cdot 10^{20}$ eV. The flux of such nuclei is expected to drop sharply at $5 \cdot 10^{19}$ eV due to the interaction with the microwave background, commonly referred to as the GZK-cutoff after its discoverers [1]. However, the number of particles known to be beyond 10^{20} eV continues to increase, with now 14 published, and a further 10 expected from new observations with HIREs and a reanalysis of the Yakutsk data [2].

There are three basic difficulties: First, we need to find a site that can either accelerate particles to such energies, or produce them outright through the decay of even higher energy particles (topological defects for example, [3]). While there are many sites which are in principle capable to produce such particles [4,5], there is only one class that has been argued to require protons at such energies in the source, namely radio galaxies with powerful jets and/or hot spots, such as M87 [6] or Cyg A. M87 has been under suspicion to be the primary source for ultra high energy cosmic rays for a long time [7,8]. For both compact and extended radiojets as well as hot spots such as in M87 the emission of synchrotron radiation [6] with a steep cutoff at frequencies about $3 \cdot 10^{14}$ Hz implies an initial turbulence injection scale of the Larmor radius of protons at 10^{21} eV. Radio galaxies of sufficient power can provide through acceleration sufficient momentum to particles to overcome losses within the space limitation and the source magnetic and radiation fields [4,6]. Gamma ray bursts are another possible class.

Secondly, there is the additional difficulty of getting these particles to us, that is, overcoming the losses in the bath of the cosmological microwave background [1]. That implies that the source should not be very much further than 20 Mpc, and that corresponds to the travel distance

for the presumably charged particles, not necessarily as short as the light travel time distance. There are only two certain candidates, the radio galaxy M87 at about 20 Mpc, and the radio galaxy NGC315 at about 80 Mpc - this latter distance is hard to accept already. Hence, we require that not much additional travel time should be spent through scattering in large scale magnetic irregularities [9].

The third difficulty is the explanation of the nearly isotropic distribution of the arrival directions of these events. The anisotropy expressed in a weak correlation with the supergalactic plane, or the detection of multiples mostly along the supergalactic plane [10,11] is a small effect. The arrival directions are best described as isotropic, at least in the accessible sky in the North. There are two extreme explanations of the isotropy, one is to argue that we have many sources contributing, even at the highest particle energies, and the other is that we have one source dominating at the maximum energies, and the particle orbits are bent. In the second option the bending should not add substantially to the travel time.

There are several ways out of these difficulties. The highest energy particles could be particles generated by topological defects [3] or from extensions of supersymmetry [12]. Alternatively, they could be cosmologically local [13,14]. If many sources contribute to the observed particle fluxes they have to be fairly common in space, as in the scenarios of accelerating such particles in the environment of quiescent black holes [15], or gamma ray bursts [16].

Here* we present the consequences of introducing a magnetic Galactic wind in analogy to the solar wind.

*inquiries to plbiermann@mpifr-bonn.mpg.de

The magnetic field of the wind bends the particle orbits without adding substantial travel time.

II. A MODEL FOR A MAGNETIC GALACTIC WIND

It has long been expected that our Galaxy has a wind [17] akin to the solar wind [18]. Recent modelling [19] shows that such winds can be quite fast, and ubiquitous.

It seems plausible that this wind is powered by the combined action of cosmic rays and magnetic fields [19] and starts in the hot phase of the interstellar medium seen in X-rays by ROSAT. This phase has a density of $3 \cdot 10^{-3}$ particles per cc, a temperature of about $4 \cdot 10^6$ K, a radial scale of about 5 kpc, and an exponential scale in z , perpendicular to the disk, of almost 2 kpc [20]. We note that the corresponding equipartition magnetic field strength is 10 microGauss.

Parker has shown [18] that in a spherical wind the poloidal component of the magnetic field becomes negligible rather quickly with radius r , the radial component decays with $1/r^2$, but the azimuthal part of the magnetic field quickly becomes dominant with $B_\phi \sim \sin \theta / r$ in polar coordinates.

An available measure of magnetic field along any line of sight is the Rotation Measure: It is proportional to the line of sight integral of the product of electron density and magnetic field component (including the sign) from us to a distant linearly polarized radio source. We verified that the magnetic field topology of the Parker model is consistent with the data [21] for a base density below that of the hot interstellar medium.

Cosmic ray driving is similar to radiation driving of winds in massive stars, and so [22] magnetic fields can lead to an increase of the momentum of the wind. For a steady wind and Parker topology the Alfvén speed in the wind becomes independent of radius r for large distances. The ultimate wind velocity is then a small multiple of the Alfvén velocity, suggesting a rather strong magnetic field. On this basis we have adopted here a magnetic field somewhat higher than in other current models [19].

The data on the sign of the azimuthal component show that in the disk of the Galaxy there are reversals but in the direction of the anti-center, the part of the sky most relevant for calculating orbits of energetic charged particles, the field points to the direction of galactic longitude about 90 degrees [23,21]. That means immediately that positively charged particles traced backwards have their origin above us, at high positive galactic latitudes.

There is one additional observational argument in favor of such a topology of the magnetic field: Krause & Beck [24] have found that the dominant magnetic field symmetry in spiral galaxies points approximately along the local spiral arms inwards towards the center. This effect is unexplained at present, but it seems nevertheless

to be observationally well established. One might expect that magnetic field sign reversals in the disk should translate into similar reversals in the wind. However, if the dominant directionality is established by the most active spiral arms, then the symmetry as found by Krause & Beck should prevail well outside the disk. We will assume this to be the case. This symmetry implies an overall current in the Galaxy possibly driven by star formation through stellar winds and cosmic rays.

If the entire disk has this symmetry, then there cannot be a sign reversal of the azimuthal component close to the disk along the direction perpendicular to the disk. This implies that the radial magnetic field component $B_r \sim 1/r^2$ is pointed inwards inside the disk and in its immediate neighborhood below and above. However, considering all of 4π the radial component has to be pointed outwards roughly over half the sky, and pointed inwards in the other half in order to satisfy the condition of a source free magnetic field. Thus, one possible and simple configuration is that B_r is pointed inwards within 30 degrees of the disk both above and below the disk, and pointed outwards within 60 degrees of both poles. Such a pattern is different from the Solar wind, where the components change sign in mid-plane.

We adopt the simplest possible model. We assume that the magnetic field in the galactic wind has a dominant azimuthal component, and ignore all other components. We assume that this azimuthal component has the same sign everywhere. This then means that the strength of this azimuthal component at the location of the Sun is the first key parameter, and the distance to which this wind extends is the second one. Since the irregularities in magnetic fields should decay into the halo, the total magnetic field strength near the Sun may be used as an approximation to the local halo wind field, and in fact one might expect a lower limit. Most measures of magnetic field underestimate its strength. Therefore we will consider for reference a model which has a field strength near the Sun of 7 microGauss [25]. The second parameter, the distance to which this wind extends, is more uncertain: Our Galaxy dominates its near environment well past our neighbor, M31, the Andromeda galaxy, and might well extend its sphere of influence to half way to M81. Therefore we will adopt as outer the halo wind radius half the distance to M81, 1.5 Mpc.

III. TRACING THE PATH BACKWARDS

To follow the particle trajectories in the Galactic halo we trace protons backwards [26] from their arrival direction at Earth. We use the 14 published cosmic ray events above 10^{20} eV, the list from Watson (included in [27]) and the new list from AGASA [28]. There is a big uncertainty with the energy estimate of the highest energy Yakutsk [2] event, which we therefore exclude from the present analysis, and hence we arrive at a final tally

of 13 events used.

Fig. 1 shows the directions of the events at that point when they leave the halo wind of our Galaxy in a polar projection. For reference we show the direction to the active galaxy M87 (Virgo A), the dominant radio galaxy in the Virgo cluster. We show the two highest energy events twice: under the assumption, (i) that they are protons, and (ii) that they are Helium-nuclei (filled black symbols). We also show the supergalactic plane as a shaded band.




FIG. 1. Directions in polar projection of the 13 highest cosmic ray events when they leave the halo of our Galaxy.

The interesting result of these model calculations is that the directions of all tracks point North. With the exception of the two events having the highest energy, all other 11 events can be traced to within less than about 20 degrees from Virgo A. Considering the uncertainty of the actual magnetic field distribution, we find then that all events are consistent with arising originally from Virgo A. Since these particles are assumed to be accelerated out of cosmic gas, about 1/10 of all particles may be Helium nuclei with the same energy per particle. If the two highest energy events are in fact He nuclei, all 13 events point within 20° of Virgo A.

If Virgo A is indeed the acceleration site of the highest energy cosmic ray events, they all require systematic bending at a ten to twenty degree level. Such bending could be easily accommodated within the plausible magnetic field strength within the supergalactic sheet from here to Virgo [29,9]. Bending by 20 degrees on a path-length of 20 Mpc implies a regular transverse magnetic field strength of order 2 nanoGauss for 10^{20} eV protons. In fact, the expected total magnetic field strength in the

supergalactic plane is significantly higher [29,21,9].

How critical are our assumptions for these results?

- The assumption of the symmetry of the magnetic field above and below the Galactic disk is important; if the magnetic field were to have opposite sign above and below the disk, then all those events that arrive below the disk, would in fact go down and point in the general direction of the Galactic South pole.
- The value of the magnetic field, here adopted as 7 microGauss, for the wind near the Sun, is a key parameter. A decrease of this value by up to $\sim 30\%$ would not change essentially the results. If the magnetic field were considerably weaker, however, of order one microGauss, then the effective pointing to a common direction of origin would be largely removed. However, it would still be sufficient to smear arrival directions, especially if its structure is regular.
- The scale of the Galactic wind here 1.5 Mpc, is not a critical parameter, since the calculations show that most of the bending happens within the first few 100 kpc.

IV. DISCUSSION AND IMPLICATIONS

This particular model can be tested in several ways:

It has been suggested repeatedly in the literature, that radio galaxies may inject a major fraction of the total energy output (of order 0.1) in energetic particles [7,30–32]. In order to determine this fraction for M87 we need a model for the magnetic field structure along the supergalactic sheet, and test it with orbit calculations.

Second, if the appearance of pairs and triplets of events [11] along the supergalactic sheet is confirmed, then clearly the magnetic field structure along the sheet may be the reason in that it gives rise to some caustics with an enhancement just along the sheet direction. Again, orbit calculations will be able to test this.

Third, the very concept of a magnetic wind, driven by cosmic rays, but with an initial magnetic field as strong as in the disk, needs to be examined more closely. The conditions to be met are a) the available energetics, b) agreement with the Rotation Measure data, c) the condition that the wind be super-Alfvénic, and d) that the wind does extend to large scales. The model for Wolf Rayet star winds [22] gives us some confidence, that this is indeed reasonable. Since this should be true for any galaxy with a high star formation rate and therefore, presumably with a high rate of cosmic ray production, this latter condition may be the easiest to test with sensitive absorption line data at high redshift, with observations of the shell of gas around these extended winds.

Fourth, the current experimental statistics cover only a fraction of the Southern sky. We would expect some asymmetry between the Northern and Southern hemispheres of our sky. How much asymmetry depends again on the more detailed magnetic field configurations, but

the data from the Auger observatory will clearly provide stringent conditions on this as on any other model.

Fifth, if the model proposed here could be confirmed, then it would constitute strong evidence that all powerful radiogalaxies produce high energy cosmic rays, and that they do this at a level of $\simeq 0.1$ of their total power output. This then implies that compact radio galaxies [12] do provide a good test bed for particle interactions, since they have a large screen of interstellar gas around the radio hot spots and jets as seen in mm-wavelength radio data [33]. There the paradigm is materialized of having a gigantic accelerator, and a beam dump. These radio galaxies may be used for particle interaction experiments in the sky. If a significant correlation in arrival direction between ultra high energy cosmic rays and this specific class of radio quasars could be confirmed [12], then properties of new particles could be constrained.

In summary, we propose here that a very simple model for a Galactic wind rather analogous to the Solar wind, may allow particle orbits at 10^{20} eV to be bent sufficiently to allow “super-GZK” particles to get here from M87, and also explains the apparent isotropy in arrival directions.

Acknowledgments. This work was started at the 1999 astrophysics summer school at the Vatican Observatory; both EJA and PLB are grateful to George Coyne, S.J. and his colleagues for inviting us and for the fruitful and inspiring atmosphere at these schools. PLB, GMT and TST would like to acknowledge also useful discussions with Rainer Beck, Pasquale Blasi, E. Boldt, Glenys Farrar, J.L. Han, Anatoly Ivanov, Randy Jokipii, Frank Jones, Hyesung Kang, Phil Kronberg, Martin Lemoine, Friedrich Meyer, Hinrich Meyer, Biman Nath, Dongsu Ryu, Michael Salomon, Günther Sigl, Alan Watson, Arnold Wolfendale and many others. We thank Phil Kronberg for a careful reading of the manuscript. Work with PLB is partially supported by a DESY-grant, GMT is partially supported by the Brazilian agencies FAPESP and CNPq, TST is supported by NASA grant NAG5-7009, and PLB and TST have jointly a grant from NATO.

-
- [1] K. Greisen. *Phys. Rev. Lett.*, **16**: 748, (1966). G. T. Zatsepin & V. A. Kuzmin. *Sov. Phys.-JETP Lett.*, **4**: 78, (1966)
 - [2] M. Takeda et al. *Phys. Rev. Lett.*, **81**: 1163, (1998). E.E. Antonov et al. *JETP Letters* **69**, 650, (1999). T. Abu-Zayyad et al. Proc. 26th ICRC, Salt Lake City, eds. D. Kieda et al. **3**: 264 (1999). A.A. Ivanov et al., ibid. **1**, 403.
 - [3] For a recent review see: P. Bhattacharjee & G. Sigl. *Physics Reports* (in press), astro-ph/9811011
 - [4] A. M. Hillas. *Ann. Rev. Astron. Astrophys.*, **22**: 425, (1984)
 - [5] P.L. Biermann. *J. Phys. G: Nucl. Part. Phys.*, **23**: 1, (1997)
 - [6] P.L. Biermann & P. Strittmatter. *Astrophys. J.*, **322**: 643, (1987)
 - [7] V.L. Ginzburg & S.I. Syrovatskii. *The origin of cosmic rays*, Pergamon Press, Oxford 1964, Russian edition 1963
 - [8] G. Cunningham et al. *Ap. J.* **236**: L71 (1980)
 - [9] D. Ryu, H. Kang & P.L. Biermann. *A&A* **335**: 19 (1998).
 - [10] T. Stanev et al. *Phys. Rev. Letters* **75**, 3056 (1995)
 - [11] N. Hayashida et al. *Phys. Rev. Lett.*, **77**: 1000, (1996). Y. Uchihori et al. In Proc. “Extremely High Energy Cosmic Rays”, Ed. M. Nagano, University of Tokyo meeting, p. 50 (1996)
 - [12] G.R. Farrar & P.L. Biermann. *Phys. Rev. Lett.*, **81**: 3579 (1998)
 - [13] V.S. Berezhinsky & M Kachelriess. *Phys. Letters B* **422**: 163 - 170 (1998), hep-ph/9709485. V.S. Berezhinsky, P. Blasi, P. & A. Vilenkin. *Phys. Rev. D* **D58**:103515 (1998), astro-ph/9803271
 - [14] T.J. Weiler. *Astropart. Phys.* **11**: 303 (1999) hep-ph/9710431.
 - [15] E. Boldt & P. Ghosh. *MNRAS* **307**: 491 (1999)
 - [16] E. Waxman & J.N. Bahcall. *Phys. Rev. Lett.*, **78**: 2292 (1997) astro-ph/9701231.
 - [17] J.A. Burke. *MNRAS* **140**: 241 (1968). H.E. Johnson & W.I. Axford. *Ap.J.* **165**: 381 (1971). W.G. Mathews, J.C. Baker. *Ap.J.* **170**: 241 (1971)
 - [18] E.N. Parker. *Ap. J.* **128**: 664 (1958).
 - [19] D. Breitschwerdt & S. Komossa. in *Astrophys. & Sp. Sc.*, Proc. “Astrophys. Dynamics”, Eds. D. Berry et al. (in press)
 - [20] S.L. Snowden et al. *Ap. J.* **485**: 125, (1997)
 - [21] P.P. Kronberg. *Rep. Prog. Phys.*, **57**: 325 (1994)
 - [22] H. Seemann & P.L. Biermann. *A&A* **327**: 273 (1997); astro-ph/9706117
 - [23] M. Simard-Normandin & P.P. Kronberg. *Nature* **279**: 115 (1979)
 - [24] F. Krause & R. Beck. *A&A* **335**: 789 (1998)
 - [25] R. Beck, A. Brandenburg, D. Moss, A. Shukurov & D. Sokoloff. *Ann. Rev. Astron. & Astrophys.* **34**: 155 (1996)
 - [26] T. Stanev. *Ap. J.* **479**: 290 (1997). G.A. Medina Tanco, E.M.de G. Dal Pino & J.E. Horvath. *Ap. J.* **492**: 200 (1998). G. Sigl & M. Lemoine. *Astropart. Phys.* **9**: 65 (1998)
 - [27] B. Wiebel-Sooth & P.L. Biermann. Landolt-Börnstein, vol. VI/3c, Springer Verlag, p. 37 (1999)
 - [28] M. Takeda, et al. *Ap. J.* **522**: 225 (1999), astro-ph/9902239.
 - [29] J.P. Vallée. *Astron. J.* **99**: 459 (1990)
 - [30] J.P. Rachen, T. Stanev, & P.L. Biermann, P.L. *Astron. & Astrophys.* **273**: 377 (1993)
 - [31] T. Enßlin et al. *Ap. J.* **477**: 560, (1997). T. Enßlin, Yiping Wang, Biman B. Nath & P.L. Biermann. *A&A* **333**: L47 (1998); astro-ph/9803105
 - [32] J.P. Leahy & N.A.B. Gizani. astro-ph/9909121
 - [33] P. P. Papadopoulos et al. *Ap. J.* in press, astro-ph/9908286